

*Application
for
United States Letters Patent*

To all whom it may concern:

Be it known that, we,
Adrian Gilbert, Rivka Riven-Kreitman, Milka Lenenberg,
Sharon Cohen-Vered and Ramon F. Joubran

have invented certain new and useful improvements in
ORAL, NASAL AND PULMONARY DOSAGE FORMULATIONS OF COPOLYMER 1

of which the following is a full, clear and exact description.

ORAL, NASAL AND PULMONARY DOSAGE FORMULATIONS OF COPOLYMER 1


5 This application is a continuation-in-part of and claims the benefit of U.S. Provisional Application No. 60/183,666, filed February 18, 2000, the contents of which are hereby incorporated by reference.

10 Throughout this application, various references are cited. These publications, in their entireties, are hereby incorporated by reference to more fully describe the state of the art to which the invention pertains.

15 **Introduction**

The present invention provides oral, nasal and pulmonary dosage formulations of Copolymer 1. COPAXONE® is the brand name for glatiramer acetate (also known as Copolymer 1). Glatiramer acetate (GA), the active ingredient of COPAXONE®, consists of the acetate salts of synthetic polypeptides, containing four naturally occurring amino acids: L-glutamic acid, L-alanine, L-tyrosine, and L-lysine with average molar fractions of [L-Glu: 0.129-0.153; L-Ala: 0.392-0.462; L-Tyr: 0.086-0.100; L-Lys: 0.300-0.374] respectively. The average molecular weight of glatiramer acetate is 4,700 - 11,000 daltons. Chemically, glatiramer acetate is designated L-glutamic acid polymer with L-alanine, L-lysine and L-tyrosine, acetate (salt). Its structural formula is:

30



35 ("Copaxone", Physician's Desk Reference, (2000), Medical Economics Co., Inc., (Montvale, NJ), 3115.) Glatiramer acetate

is also written as: poly [L-Glu¹³⁻¹⁵, L-Ala³⁹⁻⁴⁶, L-Tyr^{8.6-10}, L-Lys³⁰⁻³⁷].n CH₃COOH. Copolymer 1 is widely believed to be effective in treating a variety of immune system conditions. A common method of administering the drug is subcutaneous injection. However, such administration often results in injection site reactions such as irritation, hypersensitivity, inflammation and pain. In addition, it tends to be difficult to persuade patients to adhere to the prescribed dosing regimes. To overcome these difficulties, oral, nasal and pulmonary dosage formulations are desirable. This invention provides compositions of and processes for creating solid, semisolid and aqueous dosage forms intended for oral, nasal and pulmonary administration.

Background of the Invention

Autoimmune diseases occur when a mammal's immune system fails to recognize some of the mammal's own tissues as "self" and attacks them as "foreign". Normally, self-tolerance is developed early by developmental events within the immune system that prevent the mammal's T cell and B cells from reacting with the mammal's own tissues. Major Histocompatibility Complex (Major Histocompatibility Complex) cell surface proteins help regulate these early immune responses by binding to and presenting processed peptides to T cells.

When this self-tolerance breaks down, autoimmune diseases develop. Now, the mammal's own tissues and proteins are recognized as "antigens" and are attacked by the mammal's immune system. For example, multiple sclerosis is believed to be an autoimmune disease that occurs when the immune system attacks the myelin sheath. This sheath is thought to insulate and protect the nerves. The disease is a progressive one, characterized by demyelination, followed by neuronal loss and motor function loss. An additional example is rheumatoid arthritis ("RA"). RA is believed to be an autoimmune disease

arthritis ("RA"). RA is believed to be an autoimmune disease which involves chronic inflammation of the synovial joints and infiltration by activated T cells, macrophages and plasma cells. This inflammation is thought to lead to a progressive destruction of the articular cartilage. It is the most severe form of joint disease. The nature of the autoantigen(s) attacked in rheumatoid arthritis is poorly understood, but collagen type II is a candidate.

Like autoimmune diseases, rejection of transplanted tissue also involves a hyper-response by the immune system to an antigen. This is manifested as graft rejection in the case of organ transplantation (host-versus-graft disease, or HVGD). Another manifestation of pathological immune reactivity is graft-versus-host disease (GVHD) that occurs in approximately 30% of bone marrow recipients. Up to half of those patients who develop GVHD may succumb to this process. This high morbidity and mortality has led to continuous interest in the possibility of controlling or preventing GVHD.

There are two forms of GVHD, acute and chronic. Acute GVHD develops within the first 3 months after bone marrow transplantation and features disorders of skin, liver and gastrointestinal tract. Chronic GVHD is a multiorgan autoimmune-like disease, emerging from 3 months up to 3 years post-transplantation and shares features common to naturally occurring autoimmune disorders, like systemic lupus erythematosus (SLE) and scleroderma.

Studies on the effect of Copolymer 1 on various processes involved in the pathological course of immune rejection showed that Copolymer 1 inhibited T cell proliferation in response to host cell (Aharoni et al., Immunology Letters 58(2):79-87, 1997). Copolymer 1 treatment completely abolished cytotoxic activity toward grafts, prevented the pro-GVHD IL-2 and IFN- γ

cytokine secretion, and induced beneficial Th2 anti-inflammatory response. In view of these cumulative data, Copolymer 1 is a candidate drug for the prevention of HVGD and GVHD in humans. See WO 96/32119 and U.S. Patent No. 5,858,964.

5 Copolymer 1 has been suggested as a potential therapeutic agent for multiple sclerosis (Eur. J. Immunol. [1971] 1:242; and J. Neurol. Sci. [1977] 31:433; K.P. Johnson, *1 Neurology* 65-70 (1995); N. Engl. J. Med. [1987] 317: 408) and other immune system conditions, such as immune diseases and delayed-type
10 hyper-sensitivity conditions (WO 00/05250). This drug is a synthetic polypeptide functionally crossreactive with myelin basic protein (MBP). MBP is a natural component of the myelin sheath.

15 Copolymer 1 has been shown to suppress experimental allergic encephalomyelitis (EAE) induced by various encephalitogens, including mouse spinal cord homogenate (MSCH). MSCH encompasses all myelin antigens, such as MBP (Sela M. et al., 88 *Bull. Inst. Pasteur* 303-314 (1990), proteolipid protein (PLP) (Teitelbaum,
20 D. et al., *J. Neuroimmunol.* (1996) 64: 209-217) and myelin oligodendrocyte glycoprotein (MOG) (Ben-Nun A et al., 243 *J. Neurol.* (Suppl 1) S14-S22 (1996)) in a variety of species. EAE is an accepted model for multiple sclerosis.

25 It has also been demonstrated that Copolymer 1 is active when injected subcutaneously, intra-peritoneally, intravenously or intramuscularly (D. Teitelbaum et al., *Eur. J. Immunol.* (1971) 1:242-248; D. Teitelbaum et al., *Eur. J. Immunol.* (1973) 3:273-279). For instance, in phase III clinical trials, daily
30 subcutaneous injections of Copolymer 1 were found to slow progression of disability and reduce the relapse rate in exacerbating-remitting multiple sclerosis (K.P. Johnson, *1 Neurology* 65-70 (1995); N. Engl. J. Med. [1987] 317: 408).

35 Currently, all approved treatments of multiple sclerosis involve

subcutaneous injection of the active substance. Frequently observed injection-site reactions include irritation, hypersensitivity, inflammation and pain and even necrosis (in the case of at least one interferon β 1-b treatment) and a low level of patient compliance. Therefore, an alternative method of administration is desirable. Thus, in order to effectively treat chronic diseases such as autoimmune diseases, oral, nasal or pulmonary formulations and methods for producing such formulations are necessary.

One way to overcome the difficulties in subcutaneous injection of drugs is to create a form that can be taken orally. EP Patent 359,783 discloses the treatment of autoimmune disease by oral administration of autoantigens. In addition, it describes the oral administration of MBP for the treatment of multiple sclerosis. Oral administration of an autoantigen has been termed "oral tolerance".

PCT International Application Nos. WO 91/12816, WO 91/08760, and WO 92/06704 all depict the treatment of autoimmune diseases other than multiple sclerosis using the "oral tolerance" method with a variety of autoantigens. However, none of these references disclose the treatment of multiple sclerosis or other autoimmune diseases by the oral administration of a non-autoantigen such as Copolymer 1.

The invention also encompasses another avenue of administration that avoids the problems inherent in subcutaneous injection. This avenue is to produce a form of the drug that can be administered nasally.

Summary of the Invention

The disclosed invention provides pharmaceutical compositions useful for treating autoimmune diseases in a mammal comprising

as an active ingredient a therapeutically effective amount of Copolymer 1, and microcrystalline cellulose. Processes for the manufacture of such compositions are also disclosed.

5 **Description of the Drawings**

10 **Figure 1** shows the results of trials ascertaining prevention and treatment of EAE in Rhesus monkeys fed with Copolymer 1 (glatiramer acetate - "GA") enteric coated capsules.

15 **Figure 2** shows prevention of EAE in Rhesus monkeys fed with Copolymer 1 enteric coated capsules.

20 **Figure 3** shows the drug release profile of Copolymer 1 enteric-coated tablets.

25 **Figure 4** shows the effect of daily oral Gavage treatment with Copolymer 1 solution on rat EAE.

30 **Figure 5** shows the results of treatment of chronic-relapsing (CR)-EAE in Biozzi mice with oral Copolymer 1 solution.

35 **Figure 6** shows the results of treatment of EAE treatment in Cynomolgus monkeys by oral administration of Copolymer 1 in enteric-coated tablets.

40 **Detailed Description of the Invention**

45 This invention provides a pharmaceutical composition comprising as an active ingredient a therapeutically effective amount of Copolymer 1, and microcrystalline cellulose.

The amount of microcrystalline cellulose is at least 50 % by

weight, preferably from about 60% to about 90% by weight, more preferably at least 70 % by weight, most preferably from about 70% to about 80% by weight.

- 5 The microcrystalline cellulose may have a moisture content of up to 5.0% or a moisture content of up to 1.5%.

The pharmaceutical composition may further comprise a disintegrant. The disintegrant may be selected from the group
10 consisting of kaolin, starch, powdered sugar, sodium starch glycolate, crosscarmellose sodium, carboxymethyl cellulose, microcrystalline cellulose and sodium alginate. Preferably, the disintegrant is a pregelatinized starch. The starch may have a moisture content of up to 14%, preferably a moisture content of
15 up to 12%, more preferably a moisture content of up to 7%, most preferably a moisture content of up to 5%.

The pharmaceutical composition may further comprise a lubricant. The lubricant may be selected from the group consisting of talc,
20 sodium stearyl fumarate, magnesium stearate, calcium stearate, hydrogenated castor oil, hydrogenated soybean oil, and polyethylene glycol. Preferably, the lubricant is magnesium stearate.

- 25 The pharmaceutical composition may further comprise a protease inhibitor.

The pharmaceutical composition may further comprise an enteric coating. The enteric coating may be methacrylic ester
30 copolymer, cellulose acetate phthalate (CAP), hydroxypropyl methyl cellulose phthalate (HPMCP), carboxymethyl ethyl cellulose (CMEC), amino-alkylmethacrylate copolymer. Preferably, the enteric coating is methacrylic acid copolymer.

- 35 The pharmaceutical composition may further comprise a film

coating under the enteric coating. The film coating may be selected from the group consisting of hydroxy propyl methyl cellulose (HPMC) and poly vinyl alcohol (PVA).

- 5 The pharmaceutical composition may be in solid form. The solid form may be selected from the group consisting of a tablet, a hard gelatin capsule, a pellet and a particulate formulation.

When the pharmaceutical composition is in solid form, it may be
10 a tablet and the effective amount of Copolymer 1 is from about 0.1 mg to about 300 mg, preferably from about 5 mg to about 100 mg. In another embodiment, the effective amount of Copolymer 1 is from about 5 mg to about 50 mg. In a preferred composition, the effective amount of Copolymer 1 is about 5 mg. In another
15 preferred embodiment, the effective amount of Copolymer 1 is about 10 mg. In a further preferred composition, the effective amount of Copolymer 1 is about 50 mg. In one embodiment, the effective amount of Copolymer 1 (glatiramer acetate) is about 0.01 mg/kg to about 2 mg/kg. In a preferred composition, the
20 effective amount of Copolymer 1 (glatiramer acetate) is about 0.05 mg/kg to about 1 mg/kg.

In a preferred embodiment, the pharmaceutical composition in solid form comprises as an active ingredient a therapeutically
25 effective amount of Copolymer 1, 70%-80% by weight of microcrystalline cellulose, an enteric coating, and either 5 mg or 50 mg of Copolymer 1.

The pharmaceutical composition may further comprise a
30 pharmaceutically acceptable carrier suitable for application to mucosal linings, so as to thereby form a composition suitable for application to the mucosal linings of a subject. Preferably, the carrier is chitosan.

35 The pharmaceutical composition may further comprise a

pharmaceutically effective amount of an anti-microbial preservative. The anti-microbial preservative may be selected from the group consisting of sodium benzoate, methyl paraben, benzalkonium chloride, and propyl paraben.

5

According to this invention, the pharmaceutical composition may be in either aqueous form, or in dry powder form.

The mucosal linings to which the pharmaceutical composition may be administered may be bronchi-associated lymphoid tissue.

10

The pharmaceutical composition may be formulated for either oral administration, buccal administration, nasal administration, or pulmonary administration.

15

Also disclosed is a process for manufacturing the pharmaceutical composition, wherein the process comprises a) milling the Copolymer 1, and b) dry mixing and/or granulating the milled Copolymer 1 with at least 50% by weight of microcrystalline cellulose.

20

The process may further comprise applying a film coating or applying an enteric coating. The enteric coating may be applied using a rotating pan system.

25

Also disclosed is a method for treating an autoimmune disease in a mammal which comprises administering to the mammal the pharmaceutical composition according to this invention. The autoimmune disease is selected from the group consisting of an arthritic condition, autoimmune hemolytic anemia, autoimmune oophoritis, autoimmune uveoretinitis, Crohn's disease, chronic immune thrombocytopenic purpura, colitis, contact sensitivity disease, Graves disease, Guillain-Barre's syndrome, Hashimoto's disease, idiopathic myxedema, myasthenia gravis, psoriasis, pemphigus vulgaris, rheumatoid arthritis, GVHD or HVGD.

30

35

Preferably, the autoimmune disease is multiple sclerosis.

Thus, this invention provides a method for treating autoimmune diseases by oral administration of Copolymer 1. When Copolymer
5 1 is introduced orally, it may be mixed with other food forms and consumed in solid, semi-solid, suspension or emulsion form; and it may be mixed with pharmaceutically acceptable carriers, including water, suspending agents, emulsifying agents, adjuvants, flavor enhancers and the like. In one embodiment,
10 the oral composition is enterically-coated. Use of enteric coatings are well known in the art. Commonly known enteric coatings include Eudragit S and Eudragit L (K. Lehman, Acrylic Coatings in Controlled Release Tablet Manufacture, Manufacturing Chemist and Aerosol News, p. 39 (June 1973); K. Lehman,
15 Programmed Drug Release from Oral Program Forms, Pharma. Int., vol. ISS 3 1971, p. 34-41; Handbook of Pharmaceutical Excipients, 2nd ed.).

An additional object of the present invention is to provide
20 forms of Copolymer 1 that can be administered nasally to treat autoimmune conditions. For instance, Copolymer 1 may be administered as dry powder or metered dose of solution by inhalation, or nose-drops and nasal sprays, using appropriate formulations and metered dosing units. These formulations are
25 intended to deliver Copolymer 1 to either (i) mucosal linings of the lungs and associated airways or (ii) mucosal linings of the nasal cavities.

As contemplated, Copolymer 1 is brought into contact with those
30 lymphoid tissues in the mucosal linings which are believed to be a primary source of immune system sensitization. These mucosal linings may be found (though not necessarily exclusively) in the sinuses, trachea, bronchial passages (where they are known as the BALT or bronchi-associated lymphoid tissues) and
35 gastrointestinal linings (known as GALT or gut-associated

lymphoid tissues). Thus, the administration of Copolymer 1 is understood to include methods wherein Copolymer 1 is introduced into the body by way of ingestion or inhalation. For example, Copolymer 1 may be administered by way of the mouth through
5 feeding, through a stomach tube, by inhalation into the bronchial passages or by nasal inhalation.

The composition contemplated by the subject invention may be administered either as a simple oral solution, as an emulsion or
10 suspension formulation, as a solid oral dosage form (capsule or tablet), or even as a soft gelatin capsule. The present invention contemplates immediate release dosage forms and modified release dosage forms (including particulates, coated granules and pellets, emulsions, microemulsions and
15 encapsulation in microspheres and nanospheres).

Coated formulations may include enteric and non-enteric formulations. Enteric-coated dosage forms include the enteric-coated tablet, soft and hard gelatin capsule, pellet, particle
20 and microparticle formulations. For instance, one may employ Eudragit (methacrylic acid copolymer) or Opadry (hydroxy propyl methyl cullulose ("HPMC") or poly vinyl alcohol ("PVA")) coatings, designed to target release by pH control in the stomach or in the gut, in the duodenum, jejunum, ileum or colon.
25 Control of release may also be achieved through matrix erosion or time release formulations. Inactive ingredients can include lactose, microcrystalline cellulose, mannitol, PVP, starch, sodium starch glycolate, stearic acid, talc, hydrogenated triglycerides, polylactic acid and polyglycolic acid copolymers,
30 or other ingredients intended specifically to enhance gut absorption.

Solid oral dosage form manufacturing processes can include direct compression and tableting, dry granulation, wet
35 granulation, particulate and pellet manufacture by extrusion,

spheronisation and melt granulation. The bulk drug substance may be mixed or milled prior to blending with other excipients, or co-dissolved and sprayed in a solution in a wet granulation process. Spray drying may also be an appropriate method to achieve therapeutically important particulate formulations with or without an inactive ingredient (such as lactose). The amount of active ingredient within the composition can be in the range of 0.01 mg to 1000 mg, while the range 0.1 mg to 300 mg is likely to be optimal.

Autoimmune diseases contemplated by the present invention include either cell-mediated disease (e.g., T cell) or antibody-mediated (e.g., B cell) disorders. Such disorders can be *inter alia*, arthritic conditions, demyelinating diseases and inflammatory diseases. For example, autoimmune diseases which can be treated by the present compositions include multiple sclerosis, arthritic conditions, autoimmune hemolytic anemia, autoimmune oophoritis, autoimmune uveoretinitis, Crohn's disease, chronic immune thrombocytopenic purpura, colitis, contact sensitivity disease, Graves disease, Guillain-Barre's syndrome, Hashimoto's disease, idiopathic myxedema,, myasthenia gravis, psoriasis, pemphigus vulgaris, rheumatoid arthritis, and GVHD or HVGD. The present compositions can be used to treat one or more of these diseases.

The phrase, "arthritic condition", as used herein is a condition wherein at least one symptom of rheumatoid arthritis is observed in at least one joint of a mammal, for example in a shoulder, knee, hip, backbone or digit. Examples of arthritic conditions include "polyarthrititis", which is an arthritic condition that affects more than a single joint; "juvenile arthritis", an arthritic condition of humans under the age of 21; and Felty's syndrome, which can include the symptoms of neutropenia, splenomegaly, weight loss, anemia, lymphadenopathy, and pigment spots on the skin.

It is to be understood and expected that variations in the principles of invention herein disclosed may be made by one skilled in the art and it is intended that such modifications are to be included within the scope of the present invention.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2231

Experimental Details

EXAMPLE 1: PRODUCTION OF ENTERIC-COATED CAPSULES AND TABLETS

5 Experiment 1A

Initial pre-clinical and clinical studies were performed using enteric-coated hard gelatin capsule formulations. All formulations were tested to meet (i) assay and impurities
10 specifications, (ii) quality specifications for Copolymer 1, (iii) USP requirements for uniformity of dosage units by content uniformity and (iv) USP requirements for drug release for delayed release (enteric-coated) articles. Together, these tests were intended to ensure that each lot meets the necessary
15 requirements for identity, strength, quality and purity.

Experimental lots of enteric-coated hard gelatin capsules were prepared for initial evaluation in monkeys suffering from EAE, manufactured using co-lyophilized Copolymer 1 and mannitol,
20 sieved and hand-filled into hard gelatin capsules. The capsules were film-coated with Opadry (white) (commercial grades of hydroxy propyl methyl cellulose (HPMC) coating formulations marketed by Colorcon® (UK)) and then enteric-coated using methacrylic acid copolymer NF (Eudragit L-30 D-55, a commercial
25 grade of methacrylic acid copolymer, available as a stabilized 30% aqueous dispersion, manufactured by Rohm® (Germany)) using a Würster fluid bed dryer (Accelacota 10).

The methacrylic acid copolymer used was Eudragit L 30 D-55,
30 which is an aqueous dispersion of a methacrylic acid and acrylic acid ethyl ester. The ratio of free carbonyl groups to ester groups is 1:1. The films dissolve above pH 5.5 forming salts with alkali, thus affording coatings which are insoluble in gastric media, but soluble in the small intestine.

35

Table 1 lists the inactive ingredients employed in the hard gelatin capsules and the purpose of each component. Capsules containing 1 mg and 20 mg of active ingredient were developed.

5 TABLE 1: Excipients in Hard Gelatin Capsule Formulations

Inactive Ingredient	Pharmaceutical Function
<i>Powder-Fill</i>	
Lactose Monohydrate NF	Filler
Silicon Dioxide NF	Glidant
Pregelatinized Starch NF	Disintegrant
Magnesium Stearate NF	Lubricant
<i>Film-Coating Suspension</i>	
Opadry YS-1-7006 (clear)	Film coating
Purified Water USP	Coating suspension solvent*
<i>Enteric-Coating Suspension</i>	
Methacrylic Acid Copolymer NF (Eudragit L-30 D-55)	Enteric-coating
Talc USP	Glidant
Triethyl Citrate NF	Plasticizer
Purified Water USP	Coating suspension solvent*

*Coating solvent evaporated to dryness

i) Composition

TABLE 2

Components Strength	Composition (mg/capsule)	
	1 mg	20 mg
<i>Powder Fill</i>		
Copolymer 1	1.0	20.0
Mannitol USP	48.8	17.5
<i>Seal coat</i>		
Opadry (S-OY-7399) white	32.7	32.7
Purified Water USP	*	*
<i>Enteric Coat</i>		
Merthacrylic acid copolymer NF (Eudragit L30 D55)	20.3	20.3
Talc USP	10.3	10.3
Triethyl citrate NF	2.1	2.1
Purified Water USP	*	*

* Processing solvent evaporated to dryness.

(ii) Manufacturing Process

1. Compounding of solution of 1.2 L of Copolymer 1 and mannitol in water
2. Lyophilization in bulk in Edward's lyoflex 0.5 pilot lyophilizer
3. Sieving - mesh #20/mesh#50.
4. Capsule filing - semi-manual Feton capsule filing machine.
5. Film coating Opadry (S-OY-7399 white) in Würster fluid bed dryer.
6. Enteric coating (Eudragit L30 D55 spraying suspension) in Würster fluid bed dryer.

Rhesus monkeys were fed with the enteric coated capsules. 5

feedings before disease induction and 5 feedings after disease induction, on alternate days. Experimental allergic Encephalomyelitis (EAE) was induced by injecting 8 mg myelin basic protein (MBP) in complete Freund's adjuvant (CFA) intradermally into the leg. EAE symptoms were evaluated on the following scale:

Evaluation of EAE clinical signs:

Score	Clinical State
0	Normal neurological exam and appearance
1	Weight loss, anorexia, yawning, slow responses to stimuli, irritability or lethargy
2	Mild neurological signs, indifference, drooling, clumsiness using limbs, ataxia, tremor, altered cry and disordered gaze
3	Moderate neurological signs, blindness (pupils do not react to light), akinesia, leg weakness or paralysis
4	Severe neurological signs, semicoma, coma, quadriplegia. When the animal reached a score of 4 it was sacrificed for humane reasons
5	Death

As can be seen in Fig. 1, control animals developed EAE on day 25 post disease induction and died within 4 days from the disease.

Animals that were fed with 20 mg Copolymer 1 in enteric coated capsules did not develop any signs of disease till day 60, when it was sacrificed for histology. One animal per group was used.

The animal that was treated with 1 mg Copolymer 1 in enteric coated capsules developed some signs of disease (score 2⁺), but upon treatment with 20 mg capsules healed completely.

In a second study (Fig. 2), all monkeys treated with Copolymer 1 enteric coated capsules (1, 10, and 20 mg) were fully

protected from EAE while monkeys treated with Copolymer 1 in capsules which were not gastro-protective (opened in the stomach) or control capsule died from the disease, indicating the importance of the enteric coating in these dose levels (1 to 20 mg.).

Experiment 1B

The procedure of Experiment 1A was adapted to give a stable product under refrigerated conditions that (i) can be manufactured using standard industrial pharmaceutical processes and (ii) meets pre-defined quality specifications appropriate for initial clinical evaluation. Here, the previous experiment's process of manufacturing the powder-fill was replaced by a dry granulation process.

(i) Composition

TABLE 3

Components	composition mg/capsule		
	Placebo	20 mg	100 mg
<i>Powder Fill</i>			
Copolymer 1	0.0	20.0	100.0
Lactose monohydrate NF	322.7	36.6	183.2
Pregelatinized starch NF	37.3	6.7	33.3
Silicon Dioxide NF	9.3	1.7	8.3
Magnesium Stearate NF	3.7	0.7	3.3
<i>Seal Coat</i>			
Opadry (YS-1-7006 clear)	29.0	29.0	29.0
Purified Water USP	*	*	*
<i>Enteric Coat</i>			
Methacrylic acid copolymer NF (Eudragit L30 D55)	50.6	50.6	50.6
Talc USP	25.4	25.4	25.4
Triethyl citrate NF	8.0	8.0	8.0
Purified Water USP	*	*	*

* Processing solvent evaporated to dryness.

(ii) Manufacturing process (6kg)

1. Milling of Copolymer 1 - (fitzmill milling machine)
2. Blending with excipients - Y-cone 15 blender
3. Compression of slugs (20mm, 1g) - Killian tableting machine
4. Granulation of slugs - Frevitt granulator, 0.8 mm net
5. Blending with excipients - Y-cone 15 blender
6. Capsule filling - Bosch 400 encapsulator.
7. Coating (Accelacota-10 rotating pan machine).

The clinical formulation was administered to patients suffering from multiple sclerosis (MS) in the framework of the Phase I study. It was demonstrated that Copolymer 1 (20, 100 and 300 mg.) in enteric coated capsules was tolerated and safe in MS

patients.

In addition, oral administration of Copolymer 1 in enteric coated capsules modified the cytokine profile in multiple sclerosis patients (increase of IL-10 and decrease in IL-2) indicating a possible clinical effect of this formulation in MS patients.

Experiment 1C

10

Enteric-coated tablets (7.0 mm, round, 20 mg active ingredient) were chosen for further toxicological evaluation in monkeys. To create a stable product at controlled room temperature, Experiment 1B's procedure was modified. The lactose monohydrate filler used in the capsule formulations was replaced with microcrystalline cellulose NF (Avicel PH 102) due to the observation of long-term incompatibility with the active ingredient (a Maillard reaction occurred under accelerated storage conditions that resulted in yellowing of the granulate inside the capsule). Sodium starch glycolate NF was employed as the disintegrant; the slightly more rapid drug release brought about by sodium starch glycolate NF assisted in the toxicological evaluation in monkeys, where the gut is shorter than in humans.

25

A different opadry grade (Y-1-7000H) was used, even though this is also hydroxy propyl methyl cellulose ("HPMC")-based.

A corresponding placebo formulation was also developed.

30

The microcrystalline cellulose used was Avicel® PH 102, although Avicel® PH 101, and Avicel® PH 112, all manufactured by FMC Corporation, could be used. Microcrystalline cellulose fillers were employed in the experiments for tablet formulation. The three different kinds of microcrystalline cellulose binders mentioned differ in moisture content. Avicel® PH 101 and

Avicel® PH 102 are both high moisture content microcrystalline cellulose binders having less than or equal to 5% moisture content. Avicel® PH 112 is a low moisture content microcrystalline cellulose binder having less than or equal to 5 1.5% moisture content. The mean particle size of Avicel® ranges from 20 to 200 microns. The mean particle size of Avicel® PH 102 is 100 microns.

(i) Composition

TABLE 4

5	Ingredients	Amount (mg) /Tablet	
		0 mg	20 mg
	<i>Tablet Core</i>		
	Copolymer 1	0.0	20.0
	Microcrystalline cellulose NF	105.2	71.5
	Silicon dioxide NF	2.9	2.5
10	Sodium starch glycolate NF	5.8	5.0
	Magnesium stearate NF	1.1	1.0
	<i>Enteric Coat</i>		
15	Methacrylic acid copolymer NF (Eudragit L-30 D-55)	9.4	9.4
	Talc USP	4.7	4.7
	Triethyl citrate NF	0.9	0.9
	Purified water USP	*	*
	Total coating weight per tablet	15.0	15.0
20	Final coated tablet weight	130.0	115.0

* Processing solvent evaporated to dryness.

(ii) Manufacturing process (6 kg)

1. Milling of Copolymer 1 - (fitzmill milling machine)
- 25 2. Blending with excipients - Y-cone 15 blender
3. Compression of slugs (20 mm, 1g) - Killian tableting machine
4. Granulation of slugs - Frevitt granulator, 0.8 mm net
5. Blending with excipients - Y-cone 15 blender
- 30 6. Tableting - Killian tableting machine.
7. Coating (Accelacota-10 rotating pan machine).

Experiment 1D

Experiment 1C's procedure was adapted to create similar
5 formulations of enteric-coated tablets for clinical evaluation
in humans suffering from multiple sclerosis. Pregelatinized
starch was employed as the disintegrant. Tablet shape and size
were selected as appropriate for enteric administration and
found to be suitable for the intended pharmaceutical
10 manufacturing processes, including coating. The 5 mg active
ingredient tablets (and corresponding placebos) were
manufactured in the same shape and size as in Experiment 1C.
The size and shape of the 50 mg active ingredient tablets (and
corresponding placebos) was changed into 14.7 mm x 8.1 mm oval-
15 shaped tablets appropriate for enteric administration and the
intended pharmaceutical manufacturing processes, including
coating.

Scale-up and process qualification studies were completed for 52
20 kg lots of tablet cores for both the 5 mg and 50 mg active
ingredient. The tablet cores' composition is presented below.

(i) Composition of Tablet Cores

TABLE 5

	Ingredients	amounts of active ingredient (mg/tablet)			
		0 mg	0 mg	5 mg	50 mg
5	<i>Tablet Core</i>				
	Copolymer 1	0.0	0.0	5.0	50.0
	Microcrystalline cellulose NF (Avicel PH 102)	105.2	366.0	86.5	316.0
	Silicon dioxide NF	2.9	10.0	2.5	10.0
10	Pregelatinized starch NF	5.8	20.0	5.0	20.0
	Magnesium stearate NF	1.1	4.0	1.0	4.0
	Total	115.0	400.0	100.0	400.0
	<i>Enteric Coat</i>				
15	Methlacrylic acid copolymer NF (Eudragit L-30 D-55)	9.4	21.9	9.4	21.9
	Talc USP	4.7	11.0	4.7	11.0
	Triethyl citrate NF	0.9	2.1	0.9	2.1
	Purified water USP	*	*	*	*
	Total coating weight per tablet	15.0	35.0	15.0	35.0
20	Final coated tablet weight	130.0	435.0	115.0	435.0

* Processing solvent evaporated to dryness.

(ii) Manufacturing process (52 kg)

1. Milling of Copolymer 1 - (fitzmill milling machine)
- 25 2. Blending with excipients - Y-cone 120 blender
3. Compression of slugs (20 mm, 1g) - Killian tableting machine
4. Granulation of slugs - Frevitt granulator, 0.8 mm net
5. Blending with excipients - Y-cone 120 blender
- 30 6. Tableting - Killian tableting machine.
7. Coating sub-lots (Accelacota-10 rotating pan machine).

Accelerated stability testing (at 40°C/75% Relative Humidity) of lots with

- (i) different film-coating Opadry formulations; and
- (ii) different slug hardness values indicated that optimal formulations can be achieved by carefully controlling slug hardness and using L30 D55 Eudragit with or without Opadry film coat (or any other hydroxy propyl methyl cellulose (HPMC) formulation).

The dissolution profiles of the 5 mg and the 50 mg tablets were compared and found to be matching despite the four-fold difference in tablet weight. The dissolution profiles (in buffer at pH 6.8) are shown in Figure 3.

EXAMPLE 2: ORAL ADMINISTRATION OF COPOLYMER 1 SOLUTION

Example 2A:

Gavage Administration of Copolymer 1 in Solution to Rats

EAE was induced in Lewis rats by the injection of guinea pig spinal cord homogenate in CFA into the hind legs.

For EAE suppression, rats were fed with Copolymer 1 solution (0.1-10 mg/kg.) every day, starting from the day of EAE induction, up to day 24 (test termination).

Copolymer 1 inhibited EAE at all tested doses, with bimodal dose response pattern. The most effective doses were the lowest dose 0.1 mg/kg (see Fig. 4) and the highest dose of 10 mg/kg.

In all groups, Copolymer 1 delayed disease onset, reduced the % of sick rats, the severity of disease (as expressed by the mean score and the mean maximal score) and the disease duration.

Example 2B: Suppression of Chronic - Relapsing EAE in Biozzi
Mice by Oral Administration of Copolymer 1 in Solution

5 The chronic - relapsing EAE model represents MS better due to
its relapsing nature. It enables the demonstration of drug
effect on a disease which is on-going, similar to the situation
of treating the human disease, which is on-going.

10 Chronic - relapsing EAE was induced in female Biozzi mice by the
injection of mouse spinal cord homogenate (MSCH) in CFA,
following by re-injection of the encephalitogen one week later,
in the same manner.

15 On day 16 when the mice were already sick (see Fig. 5), they
were randomized into 4 groups and treated daily by either
phosphate buffer saline (PBS) as control or with Copolymer 1
solution (0.5 mg/kg/day, 2.5 mg/kg/day and 12.5 mg/kg/day).

20 As can be seen in Fig. 5, treatment with Copolymer 1 reduced the
EAE clinical symptoms of the second relapse in a dose-dependent
manner. The most effective dose was 12.5 mg/kg. The
suppressive effect was demonstrated in all parameters checked:
the incidence of mice suffering from second relapse, the
severity of symptoms and the duration of the relapse.

25

EXAMPLE 3: ORAL ADMINISTRATION OF COPOLYMER 1 TABLETS

Example 3A: Induction of Relapsing-Remitting EAE in Cynomolgus
Monkeys

30

A relapsing-remitting form of EAE was induced by the injection
of emulsified MBP in CFA containing 3mg/ml of Mycobacterium
Tuberculosis (MT), intradermally into right & left footpads.

35 The daily dosages employed are presented in Table 6 below.

Observation was initiated on day 8. Local reaction was observed at the site of injection, an ulcer/or bleeding ulcer caused intermittent use of the foot and /or flexion of the toes. Neurological signs appeared on day 14 following induction, as shown in Table 6.

Table 6: Daily Clinical Score

Monkey I.D.	Dose	DAYS					
		14	15	16	17	18	19-32
Female Z-920	2.5 mg	3	3	5			
Female Z-929	5 mg	0	0	3	4*		
Female Z-485	10 mg	1	2	3	5		
Male Z-523	2.5 mg	0	2	3	3/4	4*	
Male Z-1669	5 mg	0	0	0	0	0	0
Male Z-701	10 mg	0	0	0	0	0	0

* euthanized

Table 6 demonstrates that the appearance of EAE was acute, (grade 3). Two of the females died within 48 hours of onset of signs of EAE, whereas the third female was euthanized subsequent to the development of grade 4 signs (coma).

Only one male developed signs of EAE and was euthanized after it reached grade 4. The two males which did not develop signs of EAE were euthanized after 32 days.

Thus, the dose injected for the induction of EAE does not relate to the time interval between injection and appearance of signs

of EAE, or the severity of signs. The females were the first to develop the signs of EAE, and the first to die. Only a third of the males developed EAE.

5 Example 3B: Treatment of EAE in Cynomolgus Monkeys by Oral
Administration of Copolymer 1 in Enteric-Coated Tablets

Twelve Cynomolgus monkeys weighing 3 to 4 kg at the age of approximately four years were randomly divided into four
10 treatment groups. For 10 days on every alternate day (total of 5 treatments), each monkey was introduced with the tablets using tip coated forceps beyond the radix of the tongue, after which swallowing was assured. The groups were treated as follows:

- Group #1 (placebo control): four placebo tablets
15 Group #2 (5 mg Copolymer 1 per treatment): one tablet of 5 mg Copolymer 1 and 3 placebo tablets
Group #3 (10 mg Copolymer 1 per treatment): two tablets of 5 mg Copolymer 1 and 2 placebo tablets
Group #4 (20 mg Copolymer 1 per treatment): four tablets of
20 5 mg Copolymer 1.

Subsequently, following the procedure of Example 3A, EAE was induced in all monkeys by single injection with 2.5 mg of MBP per monkey. Simultaneously, each group began daily treatment
25 with the same dose of Copolymer 1 as they received prior to the induction of EAE for 30 days.

In the first study, Cop-1 treatment continued until day 52. In the second study, Cop-1 treatment continued until day 57. MBP
30 was administered in a single administration. Cop-1 was administered continuously except for several weekends in the first study, and except for Saturdays in the second study.

The monkeys were examined every Sunday to Friday, and also not
35 monitored on Saturdays, starting from day 8 after disease induction until day 52. At the end of this period, the monkeys were sacrificed and the tissue was harvested.

Local reaction was observed at the site of injection, such as an ulcer and/or bleeding ulcer, which caused intermittent use of the foot and/or flexion of the toes. EAE-related neurological
5 signs appeared between day 12 and 20 following induction, as specified in Table 7 below.

TABLE 7 Daily Clinical Score.

DOSE	0mg	0mg	0mg	5mg	5mg	5mg	10mg	10mg	10mg	20mg	20mg	20mg
	607	623	427	540	495	662	1652	1654	493	453	484	482
MONKEY ID												
5	12	0	0	0	0	0	0	0	0	0	0	2
	13	0	0	0	0	0	0	0	0	0	0	3
	14	0	0	0	0	0	0	0	0	0	0	**4
	15	0	0	0	0	0	1	0	0	0	2	4
	16	0	0	0/1	0	0	2	0	0	0/1	3	4
10	17	1	2	0/1	0	0	2	0	0	0/1	***5	4
	18	0	2	0/1	0	0	3	0	0	0	5	4
	19	0	2	0/1	0	0/1	3	2	0	0	5	4
	20	2	2	1	0/1	0/1	3	2	0/1	0	5	4
	21	2	2	1	0/1	1	3	2	0/1	0	5	4
15	22	2	2	1	0/1	0	3	2	0	0/1	5	4
	23	2	2	1	0/1	0/1	3	2	0/1	0	5	4
	24	1	0	0	0	0	2	0	0	0	5	4
	25	1	0	0	0	0	2	0	0	0	5	4
	26	0	0	0	0	0	2	0	0	0	5	4
20	27	0	0	0	0	0	2	2	0	0	5	4
	28	0	0	0	0	0	2	2	0	0	5	4
	29	0	0	0	0	0	2	2	0	0	5	4
	30	0	0	0/1	0	2	**4	0/1	0	0/1	5	4
	31											
25	32	2	0	0/1	0/1	2	4	0/1	0	0	5	4
	33	3	0	0/1	0/1	2	4	0/1	0	0	5	4
	34											
	35	3	0	1	0/1	2	4	2	0	0	5	4
	36	3	0	1	1	2	4	2	0	0	5	4
30	37											
	38	**4	0	1	1	1	4	2	0	0	5	4
	39	4	0	1	1	2	4	2	0	0	5	4
	40											
	41											
35	42	4	2	2	0	0	4	2	0	0	5	4
	43											
	44											
	45	4	2	3	0	0	4	2/3	0	0	5	4
	46	4	2	3	0/1	0	4	2	0	0	5	4
40	47	4	2	3	1	0	4	2	0	0	5	4
	48	4	2	2	1	0/1	4	2	0	0	5	4
	49	4	2	2	1	0	4	2	0	0	5	4
	50											
	51	4	2	2	1	0	4	2	0	0	5	4
45	52	4	2	2	1	0	4	2	0	0	5	4

* The monkey was sacrificed for reasons not related to neurological scoring.

35 ** The monkey was sacrificed when it reached grade 4.

*** The monkey was found dead in the cage.

All three females in the placebo group showed signs of illness on days 17-23 followed by a remission period of a week. Female No. 607 had a second acute relapse on day 32 and was sacrificed on day 38 when score 4 was reached. Females No. 623 and 427 showed severe signs of illness in the last 10 days of follow up (scores 2 and 3).

In the 5 mg Copolymer 1 treatment group, Female No. 662 showed signs of illness on day 15 and by day 18, after losing her pupillary reflex, received score 3. On day 30, the animal was sacrificed after it reached score 4. The two other females, No. 540 and 495, had a similar sequence of symptoms, with mild signs on day 18/19 to day 23 (score 1) and a relapse between day 30/32 and day 39 (score 1-2). Female No. 540 had a third relapse between day 46 to day 52 (score 1).

In the group that was treated with 10 mg Copolymer 1, Female No. 1652 showed neurological symptoms from day 19 (score 2), with a short remission period between days 24-26 and 30-33(score 0/1) and survived to the end of the follow up period. Females No. 493 and 1654 showed irregular, very mild symptoms during the entire follow up period (score fluctuated between 0 and 0/1).

In the 20 mg Copolymer 1 treatment group, Females No. 482 and 484 showed a very acute onset of signs. Female No. 482 reached score 2 on day 12, 3 on day 13 and was euthanized on day 14 (score 4).

Female No. 484 registered score 2 on day 15, reached score 3 the following day and was found dead on day 17 (score 5). Female No. 453 had mild neurological symptoms, but because of suppurative infected inguinal lymph nodes and infected bleeding wounds of the foot pad, which caused the animal severe deterioration, the animal was sacrificed on day 26.

Seven females survived to the end of the follow up period and were sacrificed on day 53. Inguinal lymph nodes and sections of the intestine were harvested for immunochemical staining.

5 The results of this example are summarized in Table 8 below.

TABLE 8: Evaluation of Clinical Manifestations

Treatment Group	Incidence	Mean Score	Mean Maximal Score	No. Relapses*	Mean Disease Onset Day
Control	3/3	1.27	3.00	3 (6)	16.66
5 mg	3/3	1.30	2.33	2 (5)	18.33
10 mg	1/3	0.49	1.33	1 (3)	23
20 mg	3/3	3.04	3.67	0 (3)	14.33

15 In summary, the course of EAE was phasic, with recurrence of symptoms and remission periods. While the placebo group showed the chronic phasic pattern of the disease, leading to deterioration with time, the 20 mg Copolymer 1 treatment group had an acute onset of symptoms with a fulminate course of the
20 disease and death. The group treated with 5 mg Copolymer 1 exhibited a pattern of disease similar to the placebo group with no obvious improvement. In the 10 mg Copolymer 1 group, two females showed irregular, very mild symptoms, while only one female showed more than very mild symptoms (score 2 with periods
25 of remission to score 0/1). All three survived through the whole study. The difference among the placebo, 10 mg Copolymer 1 treatment group and 20 mg treatment Copolymer 1 group is significant ($p < 0.0001$). However, the difference between the placebo group and the 5 mg group is not significant (n.s.) ($p =$
30 n.s.).

These results suggest that the most effective dosage is 10 mg Copolymer 1.

Example 3C: Treatment of EAE in Cynomolgus Monkeys by Oral
Administration of Copolymer 1 in Enteric-Coated Tablets

5 The procedure of Example 3B was followed with the following
exceptions - there was no 5 mg Copolymer 1 treatment group and
the monkeys were observed until day 57 after disease induction.

10 As in Example 3B, observation began on day 8 and continued until
the conclusion of the experiment, which, in this case, was day
57. Local reaction was observed at the site of injection, such
as an ulcer and/or bleeding ulcer, which caused intermittent use
of the foot and/or flexion of the toes. EAE-related
15 neurological signs appeared between day 14 and 54 as specified
in Table 9. Animals were sacrificed when they reached score 4.

TABLE 9: Daily Clinical Score.

DOSE MONKEY ID.	0mg 171	0mg 203	0mg 895	10mg 268	10mg 318	10mg 307	20mg 285	20mg 793	20mg 494
DAYS	13	0	0	0	0	0	0	0	0
	14	2	0	3	0	1	0	0	0
	15	2	0	5	0	2	0	0	0
	16	2	0	5	0	2	0	2	0
	17								
	18	3	0	5	0	3	0	4	0
	19	3	0	5	0	3	0	4	0
	20	3	0	5	0	3	0	4	0
	21	2	0	5	0	3	0	4	0
	22	2	0	5	0	3	0	4	0
	23	2	1	5	0	3	0	4	0
	24								
	25	2	1	5	0	3	0	4	0
	26	2	1	5	0	3	0	4	0
	27	3	1	5	0	3	0	4	0
	28	3	1	5	0	3	0	4	0
	29	3	1	5	0	3	0	4	0
	30	3	1	5	0	3	0	4	0
	31								
	32	4	1	5	0	3	0	4	0
	33	4	1	5	0	2	0	4	0
	34	4	1	5	0	2	0	4	0
	35	4	1	5	0	2	0	4	0
	36	4	1	5	0	2	0	4	0
	37	4	1	5	1	2	0	4	0
	38								
	39	4	1	5	1	2	1	4	0
	40	4	1	5	1	2	2	4	0
	41	4	1	5	1	2	2	4	0
	42	4	1	5	1	2	2	4	0
	43	4	1	5	1	2	2	4	0
	44	4	1	5	1	2	2	4	0
	45	4							
	46	4	1	5	1	2	2	4	0
	47	4	0	5	1	2	2	4	0
	48	4	0	5	1	2	3	4	0
	49	4	0	5	1	2	3	4	0
	50	4	0	5	1	2	3	4	0
	51	4	0	5	1	2	3	4	0
	52								
	53	4	0	5	1	2	3	4	0
	54	4	0	5	1	2	3	4	1
	55	4	0	5	1	2	3	4	1
	56	4	0	5	1	2	3	4	1
	57	4	0	5	1	2	3	4	1

2025 RELEASE

In the group that received the placebo, two females showed signs of illness on day 14 (scores 2 and 3). Female No. 895 was found dead on day 15 (score 5) and Female No. 171 was sacrificed on day 32 when score 4 was reached. Female No. 203 showed only mild signs of illness (score 1) on days 23-46.

In the 10 mg Copolymer 1 treatment group, Female No. 268 showed mild signs of illness on day 37, which continued until the end of the follow up period. Female No. 307 had a similar onset of symptoms, with mild signs on day 39 (score 1) but reached score 2 on days 40-46, and score 3 from day 46 until the end of the follow up period. Female No. 318 showed signs of illness (score 1) on day 14, reached score 2 on days 15-16, and score 3 on days 18-32, with an improvement starting on day 33 (score 2) until the end of the follow up period. Overall, she exhibited very mild symptoms during the entire follow up period.

In the group that was treated with 20 mg Copolymer 1, Female No. 285 showed a very acute onset of signs. On day 16, she reached score 2, and was euthanized two days later on day 18 (score 4). Female No. 793 had mild neurological symptoms (score 1) from day 54 until the end of follow up period. Female No. 494 showed mild signs from day 20 to 36, and reached score 2 from day 37 to 46, with an improvement (score 1) until the end of the follow up period.

A summary of the results is presented in Table 10 below.

TABLE 10: Evaluation of Clinical Manifestations

Treatment Group	Incidence	Mean Score	Mean Maximal Score	Mean Disease Onset Day
Control	3/3	2.9 ± 0.6	3.33	17
10 mg	3/3	1.3 ± 0.56	2.33	30
20 mg	3/3	1.6 ± 0.53	2.33	30

A graphic summary appears in Figure 6. Six females survived to the end of the follow up period and were sacrificed on day 57.

The phasic course of the disease was not clear as in Example 3B, with recurrence of symptoms and remission periods. Most of the animals developed acute or chronic disease. Acute onset of EAE appeared in the placebo group in two animals and in the 20 mg Copolymer 1 treatment group in one animal. In the group that was treated with 10 mg Copolymer 1, the pattern of disease was not as severe as in the other groups -- all three animals developed chronic disease and survived through the whole study. The difference between the placebo group and the 10 mg Copolymer 1 group, and between the placebo group and the 20 mg Copolymer 1 group was statistically significant ($p < 0.0001$). The difference is also significant between the 10 mg Copolymer 1 group and the 20 mg Copolymer 1 group ($p < 0.03$). From evaluation of the clinical manifestations summarized in Table 10, it is evident that Copolymer 1 treatment does not change the incidence of the disease, but mainly affects the timing of the initial manifestation of the disease and the intensity of the symptoms. This data confirms the results of Example 3B, where the most effective dosage of Copolymer 1 was 10 mg.

Example 3D: Treatment of Relapsing - Remitting (RR) MS Patients
With Copolymer 1 Enteric-Coated Tablets.

1,650 RR MS patients are recruited and randomized 1:1:1 into treatment with 5 mg Copolymer 1, 50 mg Copolymer 1 or placebo, in a framework of a phase III clinical trial. Patients are followed every 2 months for a total duration of 56 weeks.

The primary end point of the study is the reduction of relapse rate by the treatment with Copolymer 1 enteric-coated tablets.

In the secondary outcome measures, the effect of Copolymer 1 on disease activity and burden of disease are monitored using magnetic resonance imaging (MRI).

- 5 In addition, the effect on brain atrophy, magnetization transfer (MT), spectroscopy (MRS) and disability are also being monitored, in order to demonstrate the effect of oral administration of Copolymer 1 in enteric-coated tablets on disease progression.

10

DISCUSSION

- When preparing oral formulations, characteristics such as desired site of action (e.g. gut), dosage amount, physical characteristics, and chemical characteristics, must be taken
15 into consideration.

- Copolymer 1 is a non-crystalline, highly porous, lyophilized material. It is only slightly soluble, and has poor mixing and very poor flow properties. In addition, Copolymer 1 is a
20 proteinaceous material which is easily degraded by proteolytic enzymes in the gastrointestinal track. This disclosure provides Copolymer 1 formulations for oral administration which have pharmaceutical properties suitable for oral administration.

- 25 While microcrystalline cellulose has been used as a component in oral formulations, the formulations disclosed herein contain in excess of 50% by weight of microcrystalline cellulose. Such a composition, together with the disclosed milling/dry granulation manufacturing process results in oral formulations with
30 excellent flow and mixing characteristics, improved dissolution and improved stability than that which could have been expected based on the properties of Copolymer 1.

Indeed, based on the properties of Copolymer 1, it was unexpected that formulation with microcrystalline cellulose, particularly in excess of 50%, would have significantly improved pharmaceutical properties suitable for oral administration.

The advantageous properties of the disclosed formulation include that it allows for matching *in vitro* dissolution profiles of the 5 mg Copolymer 1 and the 50 mg Copolymer 1 tablets, despite a 4-fold difference in tablet weight as shown in Figure 3. Specifically, and unexpectedly, even though the 50 mg Copolymer 1 tablet is four times the weight of the 5 mg Copolymer 1 tablet, the tablets have similar dissolution profiles.